

Dickinson College

Dickinson Scholar

Faculty and Staff Publications By Year

Faculty and Staff Publications

6-2019

Legacy of Multiple Stressors: Responses of Gastropod Larvae and Juveniles to Ocean Acidification and Nutrition

Samuel Neill Bogan
Dickinson College

Julia Brumbaugh McMahon
Dickinson College

J.A. Pechenik

Anthony Pires
Dickinson College

Follow this and additional works at: https://scholar.dickinson.edu/faculty_publications



Part of the [Marine Biology Commons](#)

Recommended Citation

Bogan, S.N., J.B. McMahon, J.A. Pechenik, and A. Pires. "Legacy of Multiple Stressors: Responses of Gastropod Larvae and Juveniles to Ocean Acidification and Nutrition." *The Biological Bulletin* 236, no. 3 (2019): 159-173. <https://www.journals.uchicago.edu/doi/abs/10.1086/702993>

This article is brought to you for free and open access by Dickinson Scholar. It has been accepted for inclusion by an authorized administrator. For more information, please contact scholar@dickinson.edu.

Legacy of Multiple Stressors: Responses of Gastropod Larvae and Juveniles to Ocean Acidification and Nutrition

S. N. BOGAN^{1,2}, J. B. McMAHON^{1,3}, J. A. PECHENIK⁴, AND A. PIRES^{1,*}

¹*Department of Biology, Dickinson College, P.O. Box 1773, Carlisle, Pennsylvania 17013;* ²*Department of Ecology, Evolution, and Marine Biology, University of California, Santa Barbara, California 93106;* ³*Department of Geography and the Environment, Villanova University, 800 Lancaster Avenue, Villanova, Pennsylvania 19085;* and ⁴*Department of Biology, Tufts University, 165 Packard Avenue, Medford, Massachusetts 02145*

Abstract. Ocean acidification poses a significant threat to calcifying invertebrates by negatively influencing shell deposition and growth. An organism's performance under ocean acidification is not determined by the susceptibility of one single life-history stage, nor is it solely controlled by the direct physical consequences of ocean acidification. Shell development by one life-history stage is sometimes a function of the pH or $p\text{CO}_2$ levels experienced during earlier developmental stages. Furthermore, environmental factors such as access to nutrition can buffer organismal responses of calcifying invertebrates to ocean acidification, or they can function as a co-occurring stressor when access is low. We reared larvae and juveniles of the planktotrophic marine gastropod *Crepidula fornicata* through combined treatments of nutritional stress and low pH, and we monitored how multiple stressors endured during the larval stage affected juvenile performance. Shell growth responded non-linearly to decreasing pH, significantly declining between pH 7.6 and pH 7.5 in larvae and juveniles. Larval rearing at pH 7.5 reduced juvenile growth as a carryover effect. Larval rearing at pH 7.6 reduced subsequent juvenile growth despite the absence of a negative impact on larval growth, demonstrating a latent effect. Low larval pH magnified the impact of larval nutritional stress on competence for metamorphosis and increased carryover effects of larval nutrition on juvenile growth. Trans-life-cycle

effects of larval nutrition were thus modulated by larval exposure to ocean acidification.

Introduction

Increases in atmospheric CO_2 since the pre-industrial era are driving ocean acidification (OA) at a global scale. Hydration of dissolved CO_2 yields carbonic acid, which dissociates into free H^+ and HCO_3^- . Increased H^+ protonates dissolved CO_3^{2-} , generating additional HCO_3^- . The net effect of increased $p\text{CO}_2$ is decreased pH, decreased $[\text{CO}_3^{2-}]$, and increased $[\text{HCO}_3^-]$ (Doney *et al.*, 2009). Broadly, physiological effects of low seawater pH include acidosis, hypercapnia, and oxidative stress (Michaelidis *et al.*, 2005; Miles *et al.*, 2007; Fabry *et al.*, 2008; Tomanek *et al.*, 2011). OA also slows the biomineralization of calcium carbonate in the shells and skeletons of calcifying invertebrates such as corals, molluscs, and echinoderms (Orr *et al.*, 2005; Kurihara, 2008; Kroeker *et al.*, 2013). The relative roles of physicochemical and biological factors controlling calcification have been debated (Jokiel 2011a, b; Cyronak *et al.*, 2015; Thomsen *et al.*, 2015; Waldbusser *et al.*, 2015; Von Euw *et al.*, 2017); but the reallocation of energy between calcification, ion transport, and other physiological processes under OA has clear consequences for the growth and development of marine invertebrates (Waldbusser *et al.*, 2013; Pan *et al.*, 2015; Brunner *et al.*, 2016; Frieder *et al.*, 2017).

Organisms often exhibit stage-specific responses to OA throughout their life histories (Zippay and Hofmann, 2010; Talmage and Gobler, 2011; Ceballos-Osuna *et al.*, 2013; Dupont *et al.*, 2013; Kapsenberg and Hofmann, 2014; Small *et al.*, 2015). Investigations into the effects of OA on marine invertebrates have paid particular attention to larval development (Kurihara *et al.*, 2007; Kurihara, 2008; Miller *et al.*, 2009; Sunday *et al.*, 2011; Doropoulos *et al.*, 2012) because

Received 14 June 2018; Accepted 10 February 2019; Published online 29 April 2019.

* To whom correspondence should be addressed. Email: pires@dickinson.edu.

Abbreviations: Ω_{Ar} , saturation state of aragonite; FDR, false discovery rate; HF, high food; LF, low food; OA, ocean acidification; TA, total alkalinity; T-ISO, Tahitian *Isochrysis galbana*.

the fate of larvae is linked to settlement and population distribution (Jenkins, 2005; Johnson and Geller, 2006). While the effects of OA on larvae are important, these effects are not independent of other life-history stages. Several recent studies have documented both positive and negative effects of parental exposure to OA on the resilience of larvae to acidification stress (reviewed by Ross *et al.*, 2016). Within a generation, larval experience of OA may have consequences for later life-history stages (Byrne *et al.*, 2011; Dupont *et al.*, 2013; Gobler and Talmage, 2013). For example, in the bivalve *Ostrea lurida*, the restriction of larval shell growth by OA can persist across metamorphosis and can continue to reduce juvenile growth rate even when acidified conditions are no longer present (Hettinger *et al.*, 2012, 2013). “Carryover” effects of larval experiences on juvenile outcomes have been observed in many marine invertebrates in the contexts of a broad range of environmental stressors (Pechenik *et al.*, 1998; Pechenik, 2006, 2018).

A calcifying invertebrate may experience the costs of past acidification stress in the form of “carryover” or “latent” effects tied to their larval experience. Latent effects, as they are discussed here, are consequences of a stressor experienced during the larval period that (i) arise during later life-history stages following the recovery of larvae, or (ii) arise during later life-history stages despite the absence of a larval response (Pechenik, 2006, 2018). For example, transient nutritional deprivation of gastropod veligers can impact post-metamorphic juvenile performance even after recovery of normal larval growth rates (Pechenik *et al.*, 1996a, b, 2002). Similarly, maintaining larvae of the polychaete *Capitella* sp. at 10–12 ppt salinity for 24–48 h decreased postmetamorphic survival and growth rate, even when reduced salinity showed no effect on larval mortality (Pechenik *et al.*, 2001).

Performance under OA is also contingent on additional biotic and abiotic factors (Breitburg *et al.*, 2015; Kroeker *et al.*, 2017). A multitude of environmental stressors co-occur in coastal environments where $p\text{CO}_2$ is most variable (Reum *et al.*, 2014; Baumann *et al.*, 2015), and these co-occurring stressors can shape responses to OA in many taxa and levels of biological organization (Boyd and Hutchins, 2012; Byrne and Przeslawski, 2013; Todgham and Stillman, 2013; Ko *et al.*, 2014; Gaylord *et al.*, 2015; Griffith and Gobler, 2017; Parker *et al.*, 2017). Adequate food supply can ameliorate the effects of high $p\text{CO}_2$ and low pH in larval and juvenile bivalves (Hettinger *et al.*, 2013; Sanders *et al.*, 2013; Thomsen *et al.*, 2013; Ramajo *et al.*, 2015), as well as in corals, echinoderms, and crustaceans (Ramajo *et al.*, 2016). Conversely, low access to nutrition has been shown to exacerbate effects of OA in larval and juvenile bivalves (Thomsen *et al.*, 2013; Cole *et al.*, 2016) or to additively decrease performance in an adult bivalve (Melzner *et al.*, 2011).

Interactions between pH and nutrition may persist as carryover or latent effects across life-history stages in marine invertebrates undergoing indirect development. We tested this

hypothesis through a series of experiments using veliger larvae and juveniles of the slipper limpet, *Crepidula fornicata*. This species is a brooding planktotrophic littoral gastropod with a wide thermal tolerance (Thieltges *et al.*, 2004; Diederich and Pechenik, 2013), pH tolerance (Noisette *et al.*, 2014, 2015, 2016), and geographic distribution (Blanchard, 1997, 2009; Bohn *et al.*, 2012; JNCC, 2015). Developmental, physiological, and behavioral responses to a number of environmental factors have been studied in larvae, juveniles, and adults of this species (Pechenik, 1984; Pechenik *et al.*, 1996a, b, 2001; Diederich *et al.*, 2011; Noisette *et al.*, 2014, 2015, 2016; Padilla *et al.*, 2014; Bashevkin and Pechenik, 2015). Under optimal laboratory culture conditions, veligers of *C. fornicata* grow at about $50 \mu\text{m d}^{-1}$ with little mortality, become competent for metamorphosis at about 800- μm shell length, and are easily induced to metamorphose by exposure to 15–20 mmol L^{-1} elevated $[\text{K}^+]$ (Pechenik and Gee, 1993; Pechenik *et al.*, 1996b; Hilbish *et al.*, 1999; Padilla *et al.*, 2014). Methods for evaluating effects of nutrition on larval and juvenile development are well established in this species (Pechenik *et al.*, 2002; Pechenik and Tyrell, 2015).

We carried out four independent experiments (Table 1) in which we reared larvae and juveniles of *C. fornicata* under several different combinations of food availability and pH. We measured larval shell growth, larval survivorship, acquisition of competence for metamorphosis, and juvenile shell growth in order to understand the relationship of responses to nutrition and pH across the larva-to-juvenile transition.

Materials and Methods

Adult collection and larval culture

Two experiments (Experiments 1 and 2) were carried out at the University of Washington’s Friday Harbor Laboratories in Friday Harbor, Washington. For these experiments, brooding adults of *Crepidula fornicata* (Linnaeus, 1758) were collected during their reproductive season from the intertidal of Totten Inlet, Thurston County, Washington. Stacks of approximately 4–6 adults were housed in separate aerated 3-L glass jars containing 2 L of room temperature unfiltered seawater, which was changed daily. After adults had acclimated to the laboratory for at least 1 d, naturally hatched veligers were collected by siphoning onto a 150- μm sieve shortly after release by adults. Larvae used in Experiments 1 and 2 were derived from two separate broods. Larvae were distributed across 3 pH treatments of 8.0, 7.5, and 7.6 in Experiment 1, and 2 pH treatments of 8.0 and 7.5 in Experiment 2 (Table 1). Each pH treatment consisted of 4 replicate 800-mL jars each containing 150 larvae (Experiment 1) or 4 replicate 400-mL jars each containing 20 larvae (Experiment 2). Larvae were fed 15×10^4 cells mL^{-1} of Tahitian *Isochrysis galbana* (T-ISO), a diet that supports maximal growth rates in larvae of *C. fornicata* (e.g., Pechenik and Tyrell, 2015). T-ISO was concentrated by centrifugation and resuspended in filtered sea-

Table 1

Summary of experiments on *Crepidula fornicata*

Experiment no.	Population	Larval pH and nutrition	Larval measurements	Juvenile pH and nutrition	Age at metamorphosis (d post-hatch)	Shell length at metamorphosis ($\mu\text{m} \pm \text{SD}$)	Juvenile measurement
1	Totten Inlet, WA	8.0 HF 7.6 HF 7.5 HF	Shell growth Survivorship	–	–	–	–
2	Totten Inlet, WA	8.0 HF 7.5 HF	Shell growth Survivorship	8.0 HF 7.5 HF	12	834 \pm 104	Shell growth
3	Buzzards Bay, MA	8.0 HF 8.0 LF 7.6 HF 7.6 LF	Shell growth	8.0 HF 8.0 LF 7.6 HF 7.6 LF	19	1221 \pm 87	Shell growth
4	Buzzards Bay, MA	8.0 HF 8.0 LF 7.6 HF 7.6 LF	Shell growth Metamorphosis	8.0 HF	14 (8.0 HF) 18 (7.6 HF) 22 (7.6LF, 8.0LF)	1097 \pm 111	Shell growth

The four experiments performed during this study are described according to the geographic populations of *C. fornicata* used, the life-history stages examined, the culturing conditions applied to these stages, and what organismal processes were examined during each stage. HF, high food; LF, low food. Each juvenile growth condition included animals derived from all larval conditions in that experiment, except that in Experiment 3, only HF larvae were carried forward as juveniles.

water before addition to larval cultures, to minimize the transfer of algal growth medium. Food and culture seawater were changed every two days in Experiment 1 and daily in Experiment 2.

Two experiments (Experiments 3 and 4) were carried out at Dickinson College in Carlisle, Pennsylvania. For these experiments, adult *C. fornicata* were collected by the Marine Biological Laboratory Marine Resources Center in Woods Hole, Massachusetts, and shipped to Dickinson College during late winter. Adults were fed Shellfish Diet 1800 (Reed Mariculture, Campbell, CA) twice daily and maintained on a 14h:10h light:dark photoperiod in seawater that was bubbled with ambient air and replaced every other day. Adult incubation temperature began at 7 °C and was increased 1 °C d⁻¹ until it reached 20 °C, in order to stimulate oviposition (Pires, 2014). Upon release from brooding adults, larvae were collected as described above and distributed across larval pH and nutrition treatments, each represented by 4 replicate 800-mL culture jars each containing 150 larvae. Larval culture jars were emptied every 2 days and refilled with seawater of pH 8.0 or 7.6. High or low concentrations of T-ISO algae were added to cultures at each water change. High food (HF) cultures contained 15×10^4 cells mL⁻¹. Low food (LF) cultures contained 1×10^4 cells mL⁻¹ (e.g., Pechenik and Tyrell, 2015). Both of these experiments included all four combinations of larval pH and nutrition (8.0HF, 8.0LF, 7.6HF, and 7.6LF treatments) to enable the observation of interactive effects between larval pH and nutrition on larval and juvenile development (Table 1). Larval cultures were maintained at 20 °C on a

14h:10h photoperiod. Seawater conditions were recorded as during Experiments 1 and 2.

Seawater pH manipulation and carbonate chemistry

In two experiments (Experiments 1 and 2), seawater conditioning and carbonate chemical analyses were carried out in the Ocean Acidification Environmental Laboratory at Friday Harbor Laboratories. For Experiment 1, incoming seawater was filtered at 1 μm and equilibrated overnight at 20 °C by bubbling with ambient air (for pH 8.0) or with mixtures of CO₂ and CO₂-free air delivered by Aalborg GFC17 mass-flow controllers (Orangeburg, NY; for pH 7.5 and pH 7.6). For Experiment 2, seawater pH was dynamically conditioned at 20 °C by a feedback system that utilized a Honeywell Durafet pH electrode and Honeywell UDA2182 process controller (Charlotte, NC) to regulate injection of CO₂ and CO₂-free air to stabilize seawater pH at set points of 7.5 or 8.0 (O'Donnell *et al.*, 2013). In both Experiments 1 and 2, seawater pH was measured immediately before loading into culture jars with a Honeywell Durafet pH electrode calibrated to the total scale by the cresol purple spectrophotometric method described by Dickson *et al.*, (2007). The pH in culture jars was also measured when they were opened for water changes. In Experiment 1, headspaces of culture jars were continuously ventilated with the same gas mixtures used to condition the seawater pH treatments during the two-day intervals between culture water changes. In Experiment 2, culture jars were sealed without ventilation, and water was changed daily. Temperature and salinity

were measured with a YSI Pro Series 1030 meter (Yellow Springs, OH).

Additional studies (Experiments 3 and 4) were conducted at Dickinson College. For these experiments, seawater was collected at Buzzards Bay, Massachusetts, and filtered at 1 μm . Seawater pH was conditioned by overnight equilibration with ambient air (for pH 8.0) or with a mixture of ambient air and CO_2 delivered by Aalborg GFC17 mass-flow controllers (for pH 7.6). These gas mixtures were used to ventilate culture jar headspaces as in Experiment 1. Seawater pH was measured before and after culture water changes at two-day intervals as described for Experiment 1, but with a YSI Pro Series 1030 meter calibrated to National Institute of Standards and Technology standards. The pH measurements were converted

to total scale by recalibration of the YSI glass electrode with the cresol purple method by Dickson *et al.*, (2007). Temperature and salinity were measured as in Experiments 1 and 2.

In all experiments, seawater samples representing each treatment were fixed with mercuric chloride and were titrated to determine total alkalinity (TA) using a Mettler-Toledo DL15 automated titrator (Columbus, OH) calibrated to certified reference materials (Dickson laboratory, Scripps Institution of Oceanography). In Experiments 1–3, $p\text{CO}_2$ and Ω_{Ar} (saturation state of aragonite) were estimated based on empirical measurements of pH and TA by using CO2Sys 2.1 (Pierrot *et al.*, 2006). In Experiment 4, $p\text{CO}_2$ was also measured empirically by using an infrared CO_2 meter (CO2Meter CM-0001, Ormond Beach, FL) calibrated to an ISO-certified 1000 ppm CO_2 standard

Table 2

Physical seawater data recorded during Experiments 1–4 throughout larval and juvenile rearing of Crepidula fornicata

Experiment no.	Nominal pH (total)	Value	Actual pH (total)	Salinity (ppt)	Temperature ($^{\circ}\text{C}$)	TA ($\mu\text{mol kg}^{-1}$)	Ω_{Ar}	$p\text{CO}_2$ (μatm)	HF pH at change (total)	LF pH at change (total)
1	8.0	Mean	7.95	29.64	19.68	2097.3	1.98	490.1	7.92	–
		SD	0.02	0.15	0.66	4.3	0.10	38.3	0.03	–
		<i>n</i>	5	5	5	3	3	3	4	–
	7.6	Mean	7.63	29.86	19.88	2105.0	1.06	1103.5	7.61	–
		SD	0.03	0.15	0.13	7.7	0.10	100.3	0.03	–
		<i>n</i>	5	5	5	3	3	3	4	–
7.5	Mean	7.54	30.00	20.06	2115.3	0.89	1393.1	7.54	–	
	SD	0.02	0.20	0.30	14.6	0.07	77.7	0.03	–	
	<i>n</i>	5	5	5	3	3	3	4	–	
2	8.0	Mean	7.95	30.11	20.04	2104.6	2.04	524.2	7.88	–
		SD	0.03	0.25	0.39	6.5	0.22	69.5	0.03	–
		<i>n</i>	22	22	22	6	6	6	18	–
	7.5	Mean	7.50	30.09	20.14	2108.4	0.85	1527.3	7.51	–
		SD	0.04	0.26	0.51	12.0	0.07	87.5	0.07	–
		<i>n</i>	22	22	22	6	6	6	18	–
3	8.0	Mean	8.07	30.97	19.12	2263.7	2.63	402.9	7.98	8.03
		SD	0.04	0.19	0.57	110.7	0.16	56.7	0.04	0.05
		<i>n</i>	16	16	16	8	8	8	13	13
	7.6	Mean	7.63	30.89	19.45	2255.7	1.13	1196.9	7.55	7.58
		SD	0.03	0.35	0.75	125.8	0.06	129.2	0.04	0.05
		<i>n</i>	15	15	15	7	7	7	12	12
4	8.0	Mean	7.99	31.36	19.81	2180.8	2.29	435.3	8.15	8.05
		SD	0.02	0.49	0.27	68.8	0.21	23.3	0.10	0.09
		<i>n</i>	11	12	12	3	11	10	10	10
	7.6	Mean	7.59	31.48	20.06	2210.3	1.13	1425.2	7.78	7.59
		SD	0.03	0.26	0.39	17.5	0.13	36.4	0.09	0.05
		<i>n</i>	10	11	11	2	10	10	10	10

Values of pH are reported (from left) as the nominal treatment target values, the actual values for new seawater added to cultures, and the values recorded in high food (HF) and low food (LF) cultures immediately before regular seawater changes. Ω_{Ar} , saturation state of aragonite; TA, total alkalinity.

(Gasco, Oldsmar, FL). The chemical and physical properties of seawater used in all experiments are given in Table 2.

Induction of metamorphosis

In Experiments 2–4, larval competence for metamorphosis was assayed twice per replicate before whole cultures were exposed to a metamorphic inducer. In each assay, 10 larvae per replicate were transferred to 8 mL of seawater conditioned to the larval rearing pH containing 20 mmol L⁻¹ (Experiments 2 and 3) or 15 mmol L⁻¹ (Experiment 4) elevated KCl (Pechenik and Gee, 1993). Larvae were incubated at 20 °C in 6-well plates inside plastic boxes ventilated with the same gas mixtures used to condition seawater pH. In Experiment 2, larvae used to measure competence for metamorphosis were taken from auxiliary cultures of pH 8.0 and pH 7.6 containing 150 larvae each, so as not to deplete the sample size of larval cultures used for measuring larval growth rates. These auxiliary cultures were reared in tandem with replicates in which larval growth was measured, and they were treated with the same conditioned seawater. The frequency of metamorphosis per replicate was measured by counting the proportion of larvae that had shed their velum after 6 and 24 h of exposure to elevated KCl. Larvae tested for competence were not returned to culture. When larvae in each treatment group achieved ≥80% metamorphosis after 24 h of exposure to KCl, the remaining larvae in that treatment were exposed to KCl to produce juveniles. Metamorphosis assays performed in Experiments 2 and 3 functioned solely to determine appropriate times to produce juveniles rather than to assess the effect of larval pH or nutrition on competence, as was done in Experiment 4 by using more frequent assays.

Competence for metamorphosis was analyzed in Experiment 4 by using a full-factorial general linear model in which larval pH, larval nutrition, age at induction, and time post-induction were categorized as nominal predictor variables (Table A1). These variables were deemed nominal because of the binary nature of their treatment levels. *P*-values corresponding to individual tests within the whole-model ANOVA were adjusted to eliminate false discoveries, using Benjamini-Hochberg false discovery rate (FDR) *P*-value adjustments. Significant differences in frequencies of metamorphosis between treatment levels were evaluated using Student's *t* tests.

Juvenile culture

During Experiment 2, newly metamorphosed juveniles derived from pH 8.0 and pH 7.5 larval cultures were evenly distributed across pH 8.0 and pH 7.5 juvenile culture treatments. Juvenile treatment groups included 12 individuals, with each individual serving as a replicate. Juveniles were individually housed in 50-mL glass jars with transparent lids containing 50 mL of pH-conditioned seawater. Seawater inside juvenile jars was replaced each day with seawater of the appropriate pH and with 15×10^4 cells T-ISO mL⁻¹. Seawater condi-

tions were recorded as they were during larval culture. Juvenile cultures were incubated in a 20 °C bath and were exposed to a 12h:12h photoperiod. Juvenile growth extended for eight days.

In Experiment 3, juveniles derived from 8.0HF and 7.6HF larval treatments were evenly distributed across 4 juvenile treatments: 8.0HF, 8.0LF, 7.6HF, and 7.6LF. Juveniles derived from 8.0LF and 7.6LF larval treatments were not included because few of those larvae became competent for metamorphosis in the time available for this experiment. Experiment 4 included juveniles derived from all 4 larval treatments, but these juveniles were all cultured at pH 8.0 with HF (Table 1). Each juvenile treatment included 12 (Experiment 3) or 10 (Experiment 4) replicate individuals. These animals were individually housed in 50-mL plastic cups containing 40 mL of pH-conditioned seawater. Juvenile culture dishes of the same pH were grouped in plastic boxes ventilated with the appropriate gas mixture and were maintained at 20 °C on a 14h:10h photoperiod. Seawater and food were replaced every two days. Juvenile HF and LF concentrations were the same as those used in larval cultures. Juvenile growth extended for 8 d in Experiment 2, 14 d in Experiment 3, and 12 d in Experiment 4.

Determination of growth rates and survivorship

Growth rates of larvae and juveniles of *C. fornicata* were measured as the change of shell length over time. Individuals were nondestructively imaged using a Motic camera (Kowloon, Hong Kong) fitted to a Leica Wild M3C dissecting microscope (Leica Microsystems, Buffalo Grove, IL) and then returned to culture. Shell lengths were measured offline in ImageJ (Schneider *et al.*, 2012). Larvae were measured on their hatching day and then at four-day intervals. All individuals (*n* = 20) in each replicate culture were measured in Experiment 2. Larvae were blindly subsampled for measurement from the larger cultures used in Experiments 1, 3, and 4 (*n* = 20, 15, and 10 individuals per replicate culture, respectively). Each juvenile treatment group in Experiments 2–4 consisted of individually cultured animals, all of which were measured on the day of metamorphosis and again at three- to four-day intervals until the experiments concluded.

The survivorship of larvae was measured in Experiments 1 and 2 by counting the numbers of living larvae and empty shells within replicate cultures over time. Larval survivorship was analyzed using one-way ANOVAs with measurements that were made 10 d post-hatch. Student's *t* tests were used to perform *post hoc* comparisons of mean survivorship between treatments. All juveniles survived the duration of their experimental growth periods (Experiments 2–4); and, therefore, juvenile survivorship data are not presented. However, some juveniles developed significant epibiont growth on top of their shells during Experiment 3. This epibiont growth had the potential to obstruct movement and feeding. Four of 96 ju-

veniles were removed from culture before the completion of Experiment 3. No more than one juvenile was removed from any given treatment group.

The effects of pH and nutrition on larval shell growth were assessed throughout larval development using one-way or two-way ANOVAs to analyze shell lengths recorded at each date of measurement. Significant differences in mean shell length across treatment groups were evaluated using Student's *t* tests (Tables A2–A4).

The effects of pH and nutrition on juvenile shell growth were examined using repeated-measures analyses derived from linear mixed models. These models included the identity of each juvenile replicate as a random effect in order to account for temporal pseudoreplication. In all experiments, covariance of the “replicate identity” effect was equal to zero (Wald *P*-value = 0.0001–0.0059). Unlike larval growth data, juvenile growth measurements included sufficient replication for including a random effect within the model. For this reason, a linear mixed model was selected to analyze juvenile growth but not larval growth. Linear mixed-model parameters and results can be found in Tables A5–A7.

Results

Effects of pH and nutrition on larval performance

After 8 days of growth, larvae reared at pH 7.5 under HF began to show reduced shell lengths relative to larvae reared at pH 7.6 or pH 8.0. The shells of pH 7.5 larvae were 17% and 14% shorter than those from pH 7.6 and pH 8.0 cultures, respectively, in Experiment 1 ($P = 0.001$; Fig. 1A; Table A2) and 11% smaller than those of pH 8.0 larvae in Experiment 2 ($P = 0.015$; Fig. 1B).

Larval shell growth was less affected by rearing at pH 7.6. In Experiment 1, larvae from pH 7.6 and pH 8.0 cultures grew at the same rate during the first 8 d post-hatch (Fig. 1A). In Experiment 3, larvae reared at pH 7.6 and pH 8.0 grew at equal rates when fed LF (Fig. 1C; Table A3). When fed HF, however, larvae cultured at pH 7.6 were smaller than those cultured at pH 8.0 at 4 d ($P < 0.0001$) and 8 d ($P = 0.0064$) after hatching. By 12 d post-hatch, however, pH 7.6 larvae recovered their growth such that their shell lengths were equivalent to those cultured at pH 8.0 (Fig. 1C). Similarly, pH 8.0 and pH 7.6 larvae in Experiment 4 that were fed HF achieved equivalent shell lengths at 20 d post-hatch (Fig. 1D; Table A4). Larvae fed HF consistently grew faster than larvae fed LF at each pH level in both Experiment 3 (Table A3) and Experiment 4 (Table A3). Nutritional stress and pH stress did not appear to interactively influence the larval shell growth of *Crepidula fornicata* (Fig. 1C, D).

By 10 d post-hatch, larvae reared during Experiment 1 achieved a mean survivorship of 85.4%, and larvae reared during Experiment 2 achieved a mean survivorship of 83.6%. There was a high degree of variability in survivorship across pH treatments and experiments. Unexpectedly, in Experiment




Figure 1. Effects of seawater pH and nutrition on larval shell growth in *Crepidula fornicata*. All experiments were performed with independent broods of larvae derived from different adult broodstock. (A) Experiment 1: effects of seawater pH (7.5, 7.6, 8.0). (B) Experiment 2: effects of seawater pH (7.5, 8.0). (C) Experiment 3: combined effects of seawater pH (7.6, 8.0) and nutrition (HF, high food; LF, low food). (D) Experiment 4: combined effects of seawater pH (7.6, 8.0) and nutrition (HF, LF). In all experiments, mean shell lengths were derived from four replicate cultures per treatment. Error bars represent \pm SD.

1, larvae reared at pH 8.0 achieved 77% survivorship by 10 d post-hatch, while those reared at pH 7.6 achieved 91% and those reared at pH 7.5 achieved 88% ($F_{2,11} = 8.28$, $P = 0.0091$). In Experiment 2, 93% of pH 8.0 larvae survived by 10 d post-hatch while 72% of pH 7.5 larvae survived ($F_{1,5} = 6.87$, $P = 0.0471$), demonstrating a coupling of decreased growth rate and survivorship at pH 7.5.

Since fewer pH 8.0 larvae survived culture relative to pH 7.6 and pH 7.5 treatments during Experiment 1, it is possible that pH 8.0 larvae experienced an unknown stressor in that experiment. Such a stressor could have decreased pH 8.0 growth rates and contributed to the similarity in growth between pH 8.0 and pH 7.6 cultures during Experiment 1. However, 15% more pH 8.0 larvae survived by 10 d post-hatch in Experiment 2 compared to Experiment 1 while pH 8.0 larvae from Experiment 1 grew 18% faster than pH 8.0 larvae reared in Experiment 2. This demonstrated that survivorship and growth rate were not coupled among pH 8.0 experimental treatments.

Synergistic effects of pH and nutrition on competence for metamorphosis

Nutrition, pH, and larval age interactively affected the acquisition of competence for metamorphosis by larvae of *C. fornicata* ($F_{1,49} = 19.9$, FDR-adjusted $P = 0.0001$). At

12 d post-hatch, 55% of 8.0HF larvae metamorphosed 6 h after induction, and 95% metamorphosed after 24 h. The 7.6HF cultures were less competent by 12 d post-hatch, with only 23% of larvae metamorphosing within 6 h after induction and only 68% after 24 h. The 12-d-old LF cultures did not exhibit any metamorphosis 6 h after induction and metamorphosed at frequencies of 10% in 8.0LF and 18% in 7.6LF after 24 h (Fig. 2).

By 16 days of age, 8.0HF, 7.6HF, and 8.0LF larval cultures showed comparable frequencies of metamorphosis 6 and 24 h after they were induced to metamorphose. However, few larvae experiencing both low pH and low nutrition became competent by this age. Larvae reared at 7.6LF metamorphosed at a lower frequency than those reared at 8.0HF, 7.6HF, or 8.0LF after both 6 and 24 h of KCl exposure ($P < 0.0001$; Fig. 2).

Consequences of larval pH and nutritional stress for juvenile growth

Larval exposure to low pH influenced juvenile growth rates in three experiments (Experiments 2, 3, and 4; Fig. 3). In Experiment 2, juveniles derived from larvae that had been reared at pH 7.5 grew 33% more slowly than those derived from larvae reared at pH 8.0 ($F = 6.07, P = 0.0166$) when juveniles were cultured at pH 8.0 (Fig. 3A; Table A5). Thus, these ju-




Figure 2. Experiment 4: interactive effects of pH (7.6, 8.0) and nutrition (HF, high food; LF, low food) on competence for metamorphosis in *Crepidula fornicata*. Means are derived from 4 replicate cultures per treatment; $n = 10$ larvae per replicate. Error bars represent \pm SD. Values corresponding to different letters have significantly different means as determined by Student's t tests. Rows are separated by larval age. Columns are separated by time post-induction.




Figure 3. Effects of nutrition and pH conditioning during larval and juvenile stages on the juvenile shell growth of *Crepidula fornicata*. All experiments were performed with independent broods of larvae derived from different adult broodstock. Error bars represent \pm SD. Values measured on the same day have been minimally pushed left or right of their correct position to reduce overlap of point symbols. (A) Experiment 2: effects of larval pH (L7.5, L8.0) and juvenile pH (J7.5, J8.0) on juvenile shell growth. (B) Experiment 3: combined effects of larval pH (L7.6, L8.0), juvenile pH (J7.6, J8.0), and juvenile nutrition (HF, high food; LF, low food) on juvenile shell growth. (C) Experiment 4: combined effects of larval pH (L7.6, L8.0) and larval nutrition (HF, LF) on juvenile shell growth.

veniles incurred a carryover effect of larval acidification experience. Larval carryover effects were not apparent in juvenile *C. fornicata* reared at pH 7.5 because these cultures stopped growing 3 d post-metamorphosis regardless of their larval pH condition. Juvenile growth rates were also diminished by a main effect of juvenile pH ($F = 126.72, P < 0.0001$). *Crepidula fornicata* reared at pH 7.5 as juveniles grew 82% more slowly than pH 8.0 juveniles when individuals were derived from pH 8.0 larval cultures. Among juveniles derived from pH 7.5 larval cultures, juveniles reared at pH 7.5 grew 63% more slowly than pH 8.0 juveniles (Fig. 3A).

In Experiment 3, *C. fornicata* reared at pH 7.6 as juveniles grew at a similar rate to pH 8.0 juveniles. However, larval

culture at pH 7.6 exerted a latent effect on juvenile development that reduced post-metamorphic growth rates by 14% in juveniles fed HF and 12% in LF juveniles ($F = 29.11$, $P < 0.0001$; Fig. 3B; Table A6). Larval exposure to pH 7.6 influenced juvenile growth *via* a latent effect rather than a carryover effect, because the larvae cultured at pH 7.6 had achieved growth equivalent to their siblings cultured at pH 8.0 before they were induced to metamorphose (Fig. 1C). Moreover, the effect of larval pH on juvenile growth was influenced by juvenile diet ($F = 5.05$, $P = 0.0255$). The latent effect of larval pH 7.6 exposure was more evident among juvenile *C. fornicata* fed HF than among those fed LF (Fig. 3B).

In Experiment 4, juvenile *C. fornicata* that were fed LF as larvae grew more slowly than juveniles derived from HF larval cultures ($F = 43.86$, $P < 0.0001$; Fig. 3C; Table A7), reflecting a carryover effect similar to that caused by larval conditioning to pH 7.5. However, larval nutritional stress caused a greater carryover effect among juveniles derived from pH 7.6 larval cultures than among those derived from pH 8.0 larval cultures ($F = 5.40$, $P = 0.0214$). When larvae were reared at pH 7.6, juveniles from LF larval cultures grew 25% slower than their HF counterparts. However, when larvae were reared at pH 8.0, juveniles from LF larval cultures grew only 13% slower than their HF counterparts (Fig. 3C). Thus, the carryover effect of poor larval diet on juvenile growth was exacerbated by larval pH stress, but the direct effects of these two stressors did not interactively influence larval development. In Experiment 4, unlike in Experiments 2 and 3, larval exposure to low pH alone did not exert a negative effect on juvenile shell growth.

Discussion

Larval and juvenile growth of Crepidula fornicata only declined past a pH tipping point

The larval shell growth of *Crepidula fornicata* responded non-linearly to decreases in pH. Only after culture conditions crossed a pH threshold between 7.6 and 7.5 did mean larval growth rates significantly decrease; larval shell growth was unaffected by low-pH conditioning at pH 7.6 (Fig. 1A, C). However, larvae reared at pH 7.5 grew significantly slower than those maintained at pH 8.0 or pH 7.6 (Fig. 1A, B). Our observations on the growth of *C. fornicata* under pH 8.0, pH 7.6, and pH 7.5 are consistent with those of Kriefall *et al.* (2018), who also demonstrated subtle yet divergent transcriptomic responses of *C. fornicata* larvae to pH 7.6 and pH 7.5 conditioning, and greater transcriptional variation between pH 8.0 larvae and those in both low pH treatments. A pH tipping point was also apparent during juvenile culture. *Crepidula fornicata* did not experience reduced growth at pH 7.6, relative to pH 8.0, during its juvenile stage (Fig. 3B). Like larvae, juveniles reared at pH 7.5 grew significantly slower than those reared at pH 8.0 (Fig. 3A). Multiple experiments have thus demon-

strated that *C. fornicata* is resilient to decreased pH, incurring only minor mortality and sublethal growth effects at pH 7.5 during its larval and juvenile stages.

Non-linear calcification responses to increased $p\text{CO}_2$ or decreased pH are common among marine calcifiers (Doney *et al.*, 2009; Ries *et al.*, 2009, 2010; Waldbusser *et al.*, 2015a). Such thresholds are functions of calcium carbonate undersaturation (Doney *et al.*, 2009) or proton flux limitation (Jokiel *et al.*, 2011b), because saturation states of calcium carbonate and $[\text{H}^+]$ are tightly coupled (Cyronak *et al.*, 2015). Indeed, total pH and estimated Ω_{Ar} were correlated across experiments. The results of Experiment 1 show that larval growth began to decline between a mean pH of 7.63–7.54 and a Ω_{Ar} range of 1.06–0.89, within which aragonite began to undersaturate (Table 2). The parameter Ω_{Ar} is of interest here because aragonite is the major form of CaCO_3 that precipitates during calcification in larval and juvenile marine molluscs, including *C. fornicata* (Eyster, 1986; Falini *et al.*, 1996; Weiss *et al.*, 2002; Ries *et al.*, 2009). The decline of larval shell growth below pH 7.6 may be partially attributed to accelerated shell dissolution caused by an undersaturation of aragonite in pH 7.5 culture conditions. Likewise, Noisette *et al.* (2016) reported that net calcification rates in adult *C. fornicata* became negative at a mean pH near 7.5 in a temperature-dependent manner that was linked to Ω_{Ar} .

Larval pH stress persisted after metamorphosis as a latent or carryover effect

Rearing larvae at pH 7.5 or pH 7.6 resulted in markedly reduced rates of juvenile growth even when juveniles were reared at pH 8.0. Larval conditioning at pH 7.5 directly reduced larval growth and indirectly depressed juvenile growth after metamorphosis, thus persisting as a carryover effect (Figs. 1B, 3A). It is possible that the strength of the carryover effect of pH 7.5 conditioning was reduced by the increased mortality incurred by pH 7.5 larvae. Larvae that died during pH 7.5 exposure would likely have shown even poorer performance as juveniles had they survived through metamorphosis. Our finding that the effect of pH on growth persists across metamorphosis in *C. fornicata* as a larval carryover effect is consistent with past reports on larva-to-juvenile transitions in bivalves (Hettinger *et al.*, 2012, 2013) and echinoids (Dupont *et al.*, 2013), as well as embryo-to-larva transitions in crustaceans (Long *et al.*, 2016).

Acidification stress endured by one life-history stage has not previously been shown to affect the performance of a later life-history stage (i) when direct effects of pH are absent in both stages, or (ii) when the first stage is able to recover from acidification stress before transition to the next, as we have shown in the present work. Larvae reared at pH 7.6 recovered to pH 8.0-equivalent shell lengths before metamorphosis (Fig. 1C, D); yet juveniles that experienced pH 7.6 as larvae grew significantly more slowly than juveniles reared at

pH 8.0 as larvae, regardless of juvenile pH conditions in Experiment 3 (Fig. 3B). A latent effect caused by rearing at pH 7.6 was absent among Experiment 4 juveniles, demonstrating that trans-life-cycle effects of pH conditioning can vary across broods of *C. fornicata*. Because larval mortality was not measured in Experiments 3 and 4, it is also possible that the absence of a pH latent effect in Experiment 4 was the result of positive selection rather than inter-brood plasticity. Many taxa have been deemed resilient to OA after showing no declines in calcification or shell growth during examinations of single life-history stages (Miller *et al.*, 2009; Gazeau *et al.*, 2011; Talmage and Gobler, 2011). *Crepidula fornicata* has shown resilience to direct effects of OA during single life-history stages (Noisette *et al.*, 2015, 2016; Kriefall *et al.*, 2018), but latent pH effects from larval exposure to OA can result in additional reductions in performance.

Interactive effects of larval nutrition and larval pH before and after metamorphosis

The potential for adequate nutrition to offset physiological effects of OA has been demonstrated in multiple marine calcifiers, diet types, and life-history stages (Hettinger *et al.*, 2013b; Sanders *et al.*, 2013; Thomsen *et al.*, 2013; Pansch *et al.*, 2014; Ramajo *et al.*, 2015, 2016). Optimal nutrition did not lessen the direct effect of low pH on larval shell development in *C. fornicata* (Fig. 1C, D). Rather, low larval pH amplified negative effects of larval nutritional stress on competence for metamorphosis and juvenile shell growth.

The development and shell structure of larvae of the brooding oyster *Ostrea angasi* responded similarly to interactions between $p\text{CO}_2$ and nutrition; delayed eye development and increases in shell abnormalities in larvae with a low food supply only arose under high $p\text{CO}_2$ (Cole *et al.*, 2016). There are numerous explanations for this interaction. Mechanistically, pH stress likely impairs response pathways to nutritional deprivation that would otherwise maintain performance (Sokolova, 2013). In an energy budgeting framework, *C. fornicata* and *O. angasi* may be able to invest optimally in shell development despite nutritional stress, but that budgeted investment becomes insufficient when low pH increases energetic demands for shell development. Additionally, OA can delay the initiation of feeding in planktotrophic molluscan larvae (Gray *et al.*, 2017) and can decrease larval clearance rates (Cole *et al.*, 2016), further decreasing access to nutrition.

For juvenile *C. fornicata*, investment in shell development during low nutrition and low pH arises after the additional energetic cost of metamorphosis. A similar phenomenon (*i.e.*, interactions among larval stressors arising only after metamorphosis) was found in the congeneric species *Crepidula onyx* following larval exposure to low nutrition and hypoxic stress: larval exposure to hypoxia reduced juvenile growth, but only when larvae had poor access to food (Chiu *et al.*, 2007). In cases such as this, multiple stressors appear to dynamically in-

teract across development. This interaction is a crucial yet overlooked element of the net effect of OA on the performance and fitness of calcifying invertebrates.

The responses of growth and metamorphosis by *C. fornicata* to low pH and nutrition contrast with responses by its congener, *C. onyx*, to similar stressors (Maboloc and Chan, 2017). Larvae of *C. onyx* exhibited only minor reductions in shell length at the time of competence when reared at pH 7.38. Furthermore, *C. onyx* settled earlier, rather than later, when it was reared at low pH and fed a nutritionally poor diet (Maboloc and Chan, 2017). The differing responses between *Crepidula* species suggest differences in physiology and developmental strategy during exposure to singular and multiple stressors of OA and nutritional stress. Both species have become globally distributed across diverse environments (Blanchard, 1997; Xu, 2006) but are native to regions where low-pH events occur at distinctly different frequencies. *Crepidula onyx* is native to the Pacific Coast between Southern California and Baja, Mexico (Coe, 1935), where upwelling events reduce pH at a greater frequency and severity than in the North Atlantic, the native habitat of *C. fornicata* (Zaytsev *et al.*, 2003; Feely *et al.*, 2008). Reduction of the pre-competent phase by *C. onyx* incurring low pH and poor nutritional quality may be an adaptive response derived from its native upwelling zones. *Crepidula fornicata* has been shown to extend its pre-competent phase in response to some environmental stressors (Pechenik *et al.*, 1996b), and the delay of metamorphosis by *C. fornicata* incurring low pH and low food access indicates a developmental response to reduced pH that is distinct from that of *C. onyx*. The costs and advantages of delaying or accelerating competence in response to low pH can be comparatively examined among *Crepidula* species to better understand interactions between developmental timing and OA as they relate to performance.

Our findings suggest that the nature of interactions between multiple stressors can vary across life history and persist as trans-life-cycle effects as they are punctuated by developmental events such as metamorphosis. To accurately assess the performance of calcifying invertebrates under predicted future $p\text{CO}_2$ scenarios, a comprehensive framework that examines the relationship between multiple stressors and the legacy of those stressors should be adopted. Expanding multiple stressor studies toward *in situ* observations of environmental mosaics has proven to be a crucial step in documenting population-level consequences of OA in marine organisms (Kroeker *et al.*, 2013, 2016), but incorporating the legacy of stressors in field experiments is a challenging undertaking. Fortunately, trans-life-cycle effects of environmental stress have been successfully recorded *in situ* (Marshall *et al.*, 2003; Ng and Keough, 2003; Emler and Sadro, 2006; Chiu *et al.*, 2007; Thiyagarajan *et al.*, 2007), including carryover effects of low pH (Hettinger *et al.*, 2013). There is a growing understanding of how biotic and abiotic factors interact across life history and how they may thus impact marine calcifiers in present and future environments.

Acknowledgments

All research was supported by the National Science Foundation (CRI-OA-1416690 to Dickinson College and CRI-OA-1416846 to Tufts University). Robert Burns, Hattie Cowles, Christine Choi, Tess Dooley, Michelle Mei, Rulaiha Taylor, and Alissa Resnikoff assisted with maintenance of cultures. Rebecca Guenther, Constance Sullivan, and Michelle Herko provided technical support in the Ocean Acidification Environmental Laboratory at the University of Washington Friday Harbor Labs. Diane Cooper kindly facilitated access for field collection at the Taylor Shellfish Company. We thank Daniel Crocker for statistical advice. Critical commentary by Scott Boback, John Henson, and two anonymous reviewers materially improved the manuscript.

Literature Cited

- Bashevkin, S. M., and J. A. Pechenik. 2015. The interactive influence of temperature and salinity on larval and juvenile growth in the gastropod *Crepidula fornicata* (L.). *J. Exp. Mar. Biol. Ecol.* **470**: 78–91.
- Baumann, H., R. B. Wallace, T. Tagliaferri, and C. J. Gobler. 2015. Large natural pH, CO₂ and O₂ fluctuations in a temperate tidal salt marsh on diel, seasonal, and interannual time scales. *Estuar. Coasts* **38**: 220–231.
- Blanchard, M. 1997. Spread of the slipper limpet *Crepidula fornicata* (L. 1758) in Europe: current state and consequences. *Sci. Mar.* **61**: 109–118.
- Blanchard, M. 2009. Recent expansion of the slipper limpet population (*Crepidula fornicata*) in the Bay of Mont-Saint-Michel (Western Channel, France). *Aquat. Living Resour.* **22**: 11–19.
- Bohn, K., C. Richardson, and S. Jenkins. 2012. The invasive gastropod *Crepidula fornicata*: reproduction and recruitment in the intertidal at its northernmost range in Wales, UK, and implications for its secondary spread. *Mar. Biol.* **159**: 2091–2103.
- Boyd, P. W., and D. A. Hutchins. 2012. Understanding the responses of ocean biota to a complex matrix of cumulative anthropogenic change. *Mar. Ecol. Prog. Ser.* **470**: 125–135.
- Breitbart, D. L., J. Salisbury, J. M. Bernhard, W.-J. Cai, S. Dupont, S. C. Doney, K. J. Kroeker, L. A. Levin, W. C. Long, L. M. Milke et al. 2015. And on top of all that . . . Coping with ocean acidification in the midst of many stressors. *Oceanography* **28**: 48–61.
- Brunner, E. L., F. G. Prahl, B. Hales, and G. G. Waldbusser. 2016. A longitudinal study of Pacific oyster (*Crassostrea gigas*) larval development: isotope shifts during early shell formation reveal sub-lethal energetic stress. *Mar. Ecol. Prog. Ser.* **555**: 109–123.
- Byrne, M., and R. Przeslawski. 2013. Multistressor impacts of warming and acidification of the ocean on marine invertebrates' life histories. *Integr. Comp. Biol.* **4**: 582–596.
- Byrne, M., M. Ho, E. Wong, N. A. Soars, P. Selvakumaraswamy, H. Sheppard-Brennand, S. A. Dworjanyan, and A. R. Davis. 2011. Unshelled abalone and corrupted urchins: development of marine calcifiers in a changing ocean. *Proc. R. Soc. Biol. Sci. B* **278**: 2376–2383.
- Ceballos-Osuna, L., H. A. Carter, N. A. Miller, and J. H. Stillman. 2013. Effects of ocean acidification on early life-history stages of the porcelain crab *Petrolisthes cinctipes*. *J. Exp. Biol.* **216**: 1405–1411.
- Chiu, J. M. Y., T. Y. T. Ng, W.-X. Wang, V. Thiyagarajan, and P.-Y. Qian. 2007. Latent effects of larval food limitation on filtration rate, carbon assimilation, and growth in juvenile gastropod *Crepidula onyx*. *Mar. Ecol. Prog. Ser.* **343**: 173–182.
- Coe, W. R. 1935. Sexual phases in prosobranch mollusks of the genus *Crepidula*. *Science* **81**: 570–572.
- Cole, V. J., L. M. Parker, S. J. O'Conner, W. A. O'Conner, E. Scanes, M. Byrne, and P. Ross. 2016. Effects of multiple climate change stressors: ocean acidification interacts with warming, salinity, and low food supply on the larvae of the brooding flat oyster *Ostrea angasi*. *Mar. Biol.* **163**: 125.
- Cyronak, T., K. Schulz, and P. Jokiel. 2015. The Omega myth: what really drives lower calcification rates in an acidifying ocean. *ICES J. Mar. Sci.* **73**: 558–562.
- Dickson, A. G., C. L. Sabine, and J. R. Christian. 2007. Guide to best practices for ocean CO₂ measurements. *PICES Spec. Publ.* **3**: 64–79.
- Diederich, C. M., and J. A. Pechenik. 2013. Thermal tolerance of *Crepidula fornicata* (Gastropoda) life history stages from intertidal and subtidal subpopulations. *Mar. Ecol. Prog. Ser.* **486**: 173–187.
- Diederich, C. M., J. N. Jarrett, O. R. Chaparro, C. J. Segura, S. M. Arellano, and J. A. Pechenik. 2011. Low salinity stress experienced by larvae does not affect post-metamorphic growth or survival in three calyptraeid gastropods. *J. Exp. Mar. Biol. Ecol.* **397**: 94–105.
- Doney, S. C., V. J. Fabry, R. A. Feely, and J. A. Kleypas. 2009. Ocean acidification: the other CO₂ problem. *Annu. Rev. Mar. Sci.* **1**: 169–192.
- Doropoulos, C., S. Ward, G. Diaz-Pulido, O. Hoegh-Guldberg, and P. J. Mumby. 2012. Ocean acidification reduces coral recruitment by disrupting intimate larval-algal settlement interactions. *Ecol. Lett.* **15**: 338–346.
- Dupont, S., N. Dorey, M. Stumpp, and F. Melzner. 2013. Long-term and trans-life-cycle effects of exposure to ocean acidification in the green sea urchin *Strongylocentrotus droebachiensis*. *Mar. Biol.* **160**: 1835–1843.
- Emlet, R. B., and S. S. Sadro. 2006. Linking stages of life history: how larval quality translates into juvenile performance for an intertidal barnacle *Balanus glandula*. *Integr. Comp. Biol.* **46**: 334–346.
- Eyster, L. S. 1986. Shell inorganic composition and onset of shell mineralization during bivalve and gastropod embryogenesis. *Biol. Bull.* **170**: 211–231.
- Fabry, V., B. Seibel, R. Feely, and J. Orr. 2008. Impacts of ocean acidification on marine fauna and ecosystem processes. *ICES J. Mar. Sci.* **65**: 414–432.
- Falini, G., S. Albeck, S. Weiner, and L. Addadi. 1996. Control of aragonite or calcite polymorphism by mollusk shell macromolecules. *Science* **271**: 67–69.
- Feely, R. A., C. L. Sabine, M. Hernandez-Ayon, D. Ianson, and B. Hales. 2008. Evidence for upwelling of corrosive "acidified" water onto the continental shelf. *Science* **320**: 1490–1492.
- Frieder, C. A., S. L. Applebaum, T. F. Pan, D. Hedgecock, and D. T. Manahan. 2017. Metabolic cost of calcification in bivalve larvae under experimental ocean acidification. *ICES J. Mar. Sci.* **74**: 941–954.
- Gaylord, B., K. J. Kroeker, J. M. Sunday, K. M. Anderson, J. P. Barry, N. E. Brown, S. D. Connell, S. Dupont, K. E. Fabricius, J. M. Hall-Spencer et al. 2015. Ocean acidification through the lens of ecological theory. *Ecology* **96**: 3–15.
- Gazeau, F., J.-P. Gattuso, M. Greaves, H. Elderfield, J. Peene, C. H. R. Heip, and J. J. Middelburg. 2011. Effect of carbonate chemistry alteration on the early embryonic development of the Pacific oyster (*Crassostrea gigas*). *PLoS One* **6**: e23010.
- Gobler, C. J., and S. C. Talmage. 2013. Short- and long-term consequences of larval stage exposure to constantly and ephemerally elevated carbon dioxide for marine bivalve populations. *Biogeosciences* **10**: 2241–2253.
- Gray, M. W., C. J. Langdon, G. G. Waldbusser, B. Hales, and S. Kramer. 2017. Mechanistic understanding of ocean acidification impacts on larval feeding physiology and energy budgets of the mussel *Mytilus californianus*. *Mar. Ecol. Prog. Ser.* **563**: 81–94.
- Griffith, A. W., and C. J. Gobler. 2017. Transgenerational exposure of North Atlantic bivalves to ocean acidification renders offspring more vulnerable to low pH and additional stressors. *Sci. Rep.* **7**: 11394.
- Hettinger, A., E. Sanford, T. Hill, A. Russell, K. Sato, J. Hoey, M. Forsch, H. Page, and B. Gaylord. 2012. Persistent carry-over effects

- of planktonic exposure to ocean acidification in the Olympia oyster. *Ecology* **93**: 2758–2768.
- Hettinger, A., E. Sanford, T. Hill, E. Lenz, A. Russell, and B. Gaylord. 2013.** Larval carry-over effects from ocean acidification persist in the natural environment. *Glob. Change Biol.* **19**: 3317–3326.
- Hilbish, T. J., K. Sasada, L. S. Eyster, and J. A. Pechenik. 1999.** Relationship between rates of swimming and growth in veliger larvae: genetic variance and covariance. *J. Exp. Mar. Biol. Ecol.* **239**: 183–193.
- Jenkins, S. R. 2005.** Larval habitat selection, not larval supply, determines settlement patterns and adult distribution in two chthamalid barnacles. *J. Anim. Ecol.* **74**: 893–904.
- JNCC (Joint Nature Conservation Committee). 2015.** The Marine Habitat Classification for Britain and Ireland, Version 15.03. [Online]. Available: <http://jncc.defra.gov.uk/MarineHabitatClassification> [2019, April 16].
- Johnson, S. B., and J. B. Geller. 2006.** Larval settlement can explain the adult distribution of *Mytilus californianus* Conrad but not of *M. galloprovincialis* Lamarck or *M. trossulus* Gould in Moss Landing, central California: evidence from genetic identification of spat. *J. Exp. Mar. Biol. Ecol.* **328**: 136–145.
- Jokiel, P. L. 2011a.** Ocean acidification and control of reef coral calcification by boundary layer limitation of proton flux. *Bull. Mar. Sci.* **87**: 639–657.
- Jokiel, P. L. 2011b.** The reef coral two compartment proton flux model: a new approach relating tissue-level physiological processes to gross corallum morphology. *J. Exp. Mar. Biol. Ecol.* **409**: 1–12.
- Kapsenberg, L., and G. E. Hofmann. 2014.** Signals of resilience to ocean change: high thermal tolerance of early stage Antarctic sea urchins (*Sterechinus neumayeri*) reared under present-day and future $p\text{CO}_2$ and temperature. *Polar Biol.* **37**: 967–980.
- Ko, G. W. K., R. Dineshran, C. Campanati, V. B. S. Chan, J. Havenhand, and V. Thiagarajan. 2014.** Interactive effects of ocean acidification, elevated temperature, and reduced salinity on early-life stages of the Pacific oyster. *Environ. Sci. Technol.* **48**: 10079–10088.
- Kriefall, N. G., J. A. Pechenik, A. Pires, and S. W. Davies. 2018.** Resilience of Atlantic slippersnail *Crepidula fornicata* larvae in the face of severe coastal acidification. *Front. Mar. Sci.* **5**: 312.
- Kroeker, K. J., R. L. Kordas, R. Crim, I. E. Hendriks, L. Ramajo, G. S. Singh, C. M. Duarte, and J. P. Gattuso. 2013.** Impacts of ocean acidification on marine organisms: quantifying sensitivities and interaction with warming. *Glob. Change Biol.* **19**: 1884–1896.
- Kroeker, K. J., E. Sanford, J. M. Rose, C. A. Blanchette, F. Chan, F. P. Chavez, B. Gaylord, B. Helmuth, T. M. Hill, and G. E. Hofmann. 2016.** Interacting environmental mosaics drive geographic variation in mussel performance and predation vulnerability. *Ecol. Lett.* **19**: 771–779.
- Kroeker, K. J., R. L. Kordas, and C. D. G. Harley. 2017.** Embracing interactions in ocean acidification research: confronting multiple stressor scenarios and context dependence. *Biol. Lett.* **13**: 20160802.
- Kurihara, H. 2008.** Effects of CO_2 -driven ocean acidification on the early developmental stages of invertebrates. *Mar. Ecol. Prog. Ser.* **373**: 275–284.
- Kurihara, H., S. Kato, and A. Ishimatsu. 2007.** Effects of increased seawater $p\text{CO}_2$ on early development of the oyster *Crassostrea gigas*. *Aquat. Biol.* **1**: 91–98.
- Long, W. C., K. M. Swiney, and R. J. Foy. 2016.** Effects of high $p\text{CO}_2$ on Tanner crab reproduction and early life history. II. Carryover effects on larvae from oogenesis and embryogenesis are stronger than direct effects. *ICES J. Mar. Sci.* **73**: 836–848.
- Maboloc, E. A., and K. Chan. 2017.** Resilience of the slipper limpet *Crepidula onyx* to direct and indirect-diet effects of ocean acidification. *Sci. Rep.* **7**: 12062.
- Marshall, D. J., J. A. Pechenik, and M. J. Keough. 2003.** Larval activity levels and delayed metamorphosis affect post-larval performance in the colonial ascidian *Diplosoma listerianum*. *Mar. Biol. Prog. Ser.* **246**: 153–162.
- Melzner, F., P. Stange, K. Trübenbach, J. Thomsen, I. Casties, U. Panknin, S. Gorb, and M. Gutowska. 2011.** Food supply and seawater $p\text{CO}_2$ impact calcification and internal shell dissolution in the blue mussel *Mytilus edulis*. *PLoS One* **6**: e24223.
- Michaelidis, B., C. Ouzounis, A. Paleras, and H. Pörtner. 2005.** Effects of long-term moderate hypercapnia on acid-base balance and growth rate in marine mussels *Mytilus galloprovincialis*. *Mar. Ecol. Prog. Ser.* **293**: 109–118.
- Miles, H., S. Widdicombe, J. I. Spicer, and J. Hall-Spencer. 2007.** Effects of anthropogenic seawater acidification on acid-base balance in the sea urchin *Psammechinus miliaris*. *Mar. Pollut. Bull.* **54**: 89–96.
- Miller, A. W., A. C. Reynolds, C. Sobrino, and G. F. Riedel. 2009.** Shellfish face uncertain future in high CO_2 world: influence of acidification on oyster larvae calcification and growth in estuaries. *PLoS One* **4**: e5661.
- Ng, T. Y.-T., and M. J. Keough. 2003.** Delayed effects of larval exposure to Cu in the bryozoan *Watersipora subtorquata*. *Mar. Ecol. Prog. Ser.* **257**: 77–85.
- Noisette, F., T. Comtet, E. Legrand, F. Bordeyne, D. Davoult, and S. Martin. 2014.** Does encapsulation protect embryos from the effects of ocean acidification? The example of *Crepidula fornicata*. *PLoS One* **9**: e93021.
- Noisette, F., J. Richard, I. Le Fur, L. S. Peck, D. Davoult, and S. Martin. 2015.** Metabolic responses to temperature stress under elevated $p\text{CO}_2$ in *Crepidula fornicata*. *J. Molluscan Stud.* **81**: 238–246.
- Noisette, F., F. Bordeyne, D. Davoult, and S. Martin. 2016.** Assessing the physiological responses of the gastropod *Crepidula fornicata* to predicted ocean acidification and warming. *Limnol. Oceanogr.* **61**: 430–444.
- O'Donnell, M. J., M. N. George, and E. Carrington. 2013.** Mussel byssus attachment weakened by ocean acidification. *Nat. Clim. Change* **3**: 587–590.
- Orr, J., V. Fabry, O. Aumont, L. Bopp, S. Doney, R. A. Feely, A. Gnanadesikan, N. Gruber, A. Ishida, F. Joos et al. 2005.** Anthropogenic ocean acidification over the twenty-first century and its impact on calcifying organisms. *Nature* **437**: 681–686.
- Padilla, D. K., M. J. McCann, M. M. Glenn, A. P. Hooks, and S. E. Shumway. 2014.** Effect of food on metamorphic competence in the model system *Crepidula fornicata*. *Biol. Bull.* **227**: 242–251.
- Pan, T. F., S. L. Applebaum, and D. T. Manahan. 2015.** Experimental ocean acidification alters allocation of metabolic energy. *Proc. Natl. Acad. Sci. U.S.A.* **112**: 4696–4701.
- Pansch, C., I. Schaub, J. Havenhand, and M. Wahl. 2014.** Habitat traits and food availability determine the response of marine invertebrates to ocean acidification. *Glob. Change Biol.* **20**: 765–777.
- Parker, L. M., W. A. O'Connor, M. Byrne, R. A. Coleman, P. Virtue, M. Dove, M. Gibbs, L. Spohr, E. Scanes, and P. M. Ross. 2017.** Adult exposure to ocean acidification is maladaptive for larvae of the Sydney rock oyster *Saccostrea glomerata* in the presence of multiple stressors. *Biol. Lett.* **13**: 20160798.
- Pechenik, J. A. 1984.** The relationship between temperature, growth rate, and duration of planktonic life for larvae of the gastropod *Crepidula fornicata* (L.). *J. Exp. Mar. Biol. Ecol.* **74**: 241–257.
- Pechenik, J. A. 2006.** Larval experience and latent effects: Metamorphosis is not a new beginning. *Integr. Comp. Biol.* **46**: 323–333.
- Pechenik, J. A. 2018.** Latent effects: surprising consequences of embryonic and larval experience on life after metamorphosis. Pp. 208–225 in *Evolutionary Ecology of Marine Invertebrate Larvae*, T. J. Carrier, A. M. Rietzel, and A. Heyland, eds. Oxford University Press, Oxford.
- Pechenik, J. A., and C. C. Gee. 1993.** Onset of metamorphic competence in larvae of the gastropod *Crepidula fornicata* (L.), judged by a natural and artificial cue. *J. Exp. Mar. Biol. Ecol.* **167**: 59–72.
- Pechenik, J. A., and A. Tyrell. 2015.** Larval diet alters larval growth rates and post-metamorphic performance in the marine gastropod *Crepidula fornicata*. *Mar. Biol.* **162**: 1597–1610.

- Pechenik, J. A., M. S. Estrella, and K. Hammer. 1996a.** Food limitation stimulates metamorphosis of competent larvae and alters postmetamorphic growth rate in the marine prosobranch gastropod *Crepidula fornicata*. *Mar. Biol.* **127**: 267–275.
- Pechenik, J. A., K. Hammer, and C. Weise. 1996b.** The effect of starvation on acquisition of competence and post-metamorphic performance in the marine prosobranch gastropod *Crepidula fornicata* (L.). *J. Exp. Mar. Biol. Ecol.* **199**: 137–152.
- Pechenik, J. A., D. Wendt, and J. Jarrett. 1998.** Metamorphosis is not a new beginning. *BioScience* **48**: 901–910.
- Pechenik, J. A., T. Gleason, D. Daniels, and D. Champlin. 2001.** Influence of larval exposure to salinity and cadmium stress on juvenile performance of two marine invertebrates (*Capitella* sp. and *Crepidula fornicata*). *J. Exp. Mar. Biol. Ecol.* **264**: 101–114.
- Pechenik, J. A., J. Jarrett, and J. Rooney. 2002.** Relationships between larval nutritional experience, larval growth rates, juvenile growth rates, and juvenile feeding rates in the prosobranch gastropod *Crepidula fornicata*. *J. Exp. Mar. Biol. Ecol.* **280**: 63–78.
- Pierrot, D., E. Lewis, and D. W. R. Wallace. 2006.** MS Excel Program Developed for CO₂ System Calculations. ORNL/CDIAC-105a. Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, U.S. Department of Energy, Oak Ridge, TN.
- Pires, A. 2014.** Artificial seawater culture of the gastropod *Crepidula fornicata* for studies of larval settlement and metamorphosis. Pp. 35–44 in *Developmental Biology of the Sea Urchin and Other Marine Invertebrates: Methods and Protocols*, S. Stricker and D. Carroll, eds. Series: Methods in Molecular Biology, 1128. Humana Press, New York.
- Ramajo, L., N. Marba, L. Prado, S. Peron, M. A. Lardies, A. B. Rodriguez-Navarro, C. A. Vargas, N. A. Lagos, and C. M. Duarte. 2015.** Biomineralization changes with food supply confer juvenile scallops (*Argopecten purpuratus*) resistance to ocean acidification. *Glob. Change Biol.* **22**: 2025–2037.
- Ramajo, L., E. Pérez-León, I. E. Hendriks, N. Marbà, D. Krause-Jensen, M. K. Sejr, M. E. Blicher, N. A. Lagos, Y. S. Olsen, and C. M. Duarte. 2016.** Food supply confers calcifiers resistance to ocean acidification. *Sci. Rep.* **6**: 19374.
- Reum, J. C. P., S. R. Alin, R. A. Feely, J. Newton, M. Warner, and P. McElhany. 2014.** Seasonal carbonate chemistry covariation with temperature, oxygen, and salinity in a fjord estuary: implications for the design of ocean acidification experiments. *PLoS One* **9**: e89619.
- Ries, J. B., A. L. Cohen, and D. C. McCorkle. 2009.** Marine calcifiers exhibit mixed responses to CO₂-induced ocean acidification. *Geology* **37**: 1131–1134.
- Ries, J. B., A. L. Cohen, and D. C. McCorkle. 2010.** A nonlinear calcification response to CO₂-induced ocean acidification by the coral *Oculina arbuscula*. *Coral Reefs* **29**: 661674.
- Ross, P. M., L. Parker, and M. Byrne. 2016.** Transgenerational responses of molluscs and echinoderms to changing ocean conditions. *ICES J. Mar. Sci.* **73**: 537–549.
- Sanders, M. B., T. P. Bean, T. H. Hutchinson, and W. J. F. Le Quesne. 2013.** Juvenile king scallop, *Pecten maximus*, is potentially tolerant to low levels of ocean acidification when food is unrestricted. *PLoS One* **8**: e74118.
- Schneider, C. A., W. S. Rasband, and K. W. Eliceiri. 2012.** NIH Image to ImageJ: 25 years of image analysis. *Nat. Methods* **9**: 671–675.
- Small, D. P., P. Calosi, D. Boothroyd, S. Widdicombe, and J. I. Spicer. 2015.** Stage-specific changes in physiological and life-history responses to elevated temperature and pCO₂ during the larval development of the European lobster *Homarus gammarus* (L.). *Physiol. Biochem. Zool.* **88**: 494–507.
- Sokolova, I. M. 2013.** Energy-limited tolerance to stress as a conceptual framework to integrate the effects of multiple stressors. *Integr. Biol.* **53**: 597–608.
- Sunday, J., R. Crim, C. Harley, and M. Hart. 2011.** Quantifying rates of evolutionary adaptation in response to ocean acidification. *PLoS One* **6**: e22881.
- Talmage, S., and C. Gobler. 2011.** Effects of elevated temperature and carbon dioxide on the growth and survival of larvae and juveniles of three species of northwest Atlantic bivalves. *PLoS One* **6**: e26941.
- Thieltges, D. W., M. Strasser, J. E. E. van Beusekom, and K. Reise. 2004.** Too cold to prosper—winter mortality prevents population increase of the introduced American slipper limpet *Crepidula fornicata* in northern Europe. *J. Exp. Mar. Biol. Ecol.* **311**: 375–391.
- Thiyagarajan, V., J. A. Pechenik, L. A. Gosselin, and P. Y. Qian. 2007.** Juvenile growth in barnacles: combined effect of delayed metamorphosis and sublethal exposure of cyprids to low-salinity stress. *Mar. Ecol. Prog. Ser.* **344**: 173–184.
- Thomsen, J., I. Casties, C. Pansch, A. Körtzinger, and F. Melzner. 2013.** Food availability outweighs ocean acidification in juvenile *Mytilus edulis*: laboratory and field experiments. *Glob. Change Biol.* **4**: 1017–1027.
- Thomsen, J., K. Haynert, K. M. Wegner, and F. Melzner. 2015.** Impact of seawater carbonate chemistry on the calcification of marine bivalves. *Biogeosciences* **12**: 4209–4220.
- Todgham, A. E., and J. H. Stillman. 2013.** Physiological responses to shifts in multiple environmental stressors: relevance in a changing world. *Integr. Comp. Biol.* **53**: 539–544.
- Tomanek, L., M. Zuzow, A. Ivanina, E. Beniash, and I. Sokolova. 2011.** Proteomic response to elevated pCO₂ level in eastern oysters, *Crassostrea virginica*: evidence for oxidative stress. *J. Exp. Biol.* **214**: 1836–1844.
- Von Euw, S., Q. Zhang, V. Manichev, N. Murali, J. Gross, L. C. Feldman, T. Gustafsson, C. Flach, R. Mendelshon, and P. G. Falkowski. 2017.** Biological control of aragonite formation in stony corals. *Science* **356**: 933–938.
- Waldbusser, G. G., E. Brunner, B. Hales, B. Ales, C. Langdon, and F. Prahl. 2013.** A developmental and energetic basis linking larval oyster shell formation to acidification sensitivity. *Geophys. Res. Lett.* **40**: 2171–2176.
- Waldbusser, G. G., B. Hales, C. J. Langdon, B. A. Hales, P. Schrader, E. L. Brunner, M. W. Gray, C. A. Miller, and I. Gimenez. 2015.** Saturation-state sensitivity of marine bivalve larvae to ocean acidification. *Nat. Clim. Change* **5**: 273–280.
- Weiss, I., N. Tuross, L. Addadi, and S. Weiner. 2002.** Mollusc larval shell formation: Amorphous calcium carbonate is a precursor phase for aragonite. *J. Exp. Zool.* **293**: 478–491.
- Xu, H., S. Qiang, Z. Han, J. Guo, Z. Huang, H. Sun, S. He, H. Ding, H. Wu, and F. Wan. 2006.** The status and causes of alien species invasion in China. *Biodivers. Conserv.* **15**: 2893–2904.
- Zaytsev, O., R. Cervantes-Duarte, O. Montante, and A. Gellegos-Garcia. 2003.** Coastal upwelling activity on the Pacific shelf of the Baja California border. *J. Oceanogr.* **49**: 489–502.
- Zippay, M. L., and G. E. Hofmann. 2010.** Effect of pH on gene expression and thermal tolerance of early life history stages of red abalone. *J. Shellfish Res.* **29**: 429–439.

Appendix

Table A1

Experiment 4: general linear model parameters and results for mean frequencies of larval metamorphosis of *Crepidula fornicata*

Fixed effect	df (test, error)	F-ratio	FDR <i>P</i> -value	Sum of squares
Whole model	14, 49	29.78	< 0.0001	6.94
Lack of fit	1, 48	0.04	0.8487	0.01
pH	1, 49	23.01	< 0.0001	0.39
Nutrition	1, 49	155.52	< 0.0001	2.64
Time post-induction	1, 49	107.34	< 0.0001	1.82
Age	1, 49	74.53	< 0.0001	1.27
pH × nutrition	1, 49	0.59	0.5625	0.01
pH × time post-induction	1, 49	0.33	0.6576	0.01
pH × age	1, 49	0.59	0.5625	0.01
Nutrition × time post-induction	1, 49	4.45	0.0667	0.08
Nutrition × age	1, 49	14.72	0.0007	0.25
Time post-induction × age	1, 49	2.98	0.1360	0.05
pH × nutrition × age	1, 49	19.47	0.0001	0.33
pH × nutrition × time post-induction	1, 49	0.00	1	0
pH × age × time post-induction	1, 49	0.14	0.7539	0.01
Nutrition × age × time post-induction	1, 49	5.30	0.0485	0.09

Significant effects (Benjamini-Hochberg false discovery rate [FDR]-adjusted $P < 0.05$) are indicated in bold.

Table A2

Experiment 1: multiple one-way ANOVA results assessing the effect of pH on larval shell length of *Crepidula fornicata* at different times post-hatch

Larval age (d)	<i>F</i> -ratio	<i>P</i> -value	<i>P</i> -value		
			8.0 vs. 7.6	8.0 vs. 7.5	7.6 vs. 7.5
4	4.08	0.0548	na	na	na
8	26.77	0.0002	0.1968	0.0004	< 0.0001

Significant differences ($P < 0.05$) demonstrated by Student's *t* tests are indicated in bold. "na" indicates *post hoc* comparisons that could not be made because of lack of a significant corresponding effect.

Table A3

Experiment 3: multiple two-way ANOVA results assessing the effect of pH and nutrition on larval shell length of *Crepidula fornicata* at different times post-hatch

Larval age (d)	Fixed effect	F-ratio	P-value	P-value					
				8.0HF vs. 8.0LF	7.6HF vs. 7.6LF	8.0HF vs. 7.6LF	8.0LF vs. 7.6HF	8.0HF vs. 7.6HF	8.0LF vs. 7.6LF
4	Nutrition	223.71	<0.0001	<0.0001					
	pH	59.07	<0.0001	–	–	<0.0001			
	Interaction	26.61	0.0002	<0.0001	<0.0001	0.0002	<0.0001	<0.0001	0.0992
8	Nutrition	860.40	<0.0001	<0.0001					
	pH	0.94	0.0037	–	–		0.0037		
	Interaction	0.02	0.0419	<0.0001	<0.0001	<0.0001	<0.0001	0.0014	0.3717
12	Nutrition	860.40	<0.0001	<0.0001					
	pH	0.94	0.3522	–	–		na		
	Interaction	0.02	0.8821	na	na	na	na	na	na

Significant differences ($P < 0.05$) demonstrated by Student's *t* tests are indicated in bold. *P*-values in merged cells that combine high food (HF) vs. low food (LF) or pH 8.0 vs. pH 7.6 columns correspond to *post hoc* comparisons of means for all HF or LF or pH 8.0 or pH 7.6 data. "na" indicates *post hoc* comparisons that could not be made because of lack of a significant corresponding effect.

Table A4

Experiment 4: multiple two-way ANOVA results assessing the effect of pH and nutrition on larval shell length of *Crepidula fornicata* at different times post-hatch

Larval age (d)	Fixed effect	F-ratio	P-value	P-value					
				8.0HF vs. 8.0LF	7.6HF vs. 7.6LF	8.0HF vs. 7.6LF	8.0LF vs. 7.6HF	8.0HF vs. 7.6HF	8.0LF vs. 7.6LF
4	Nutrition	71.08	<0.0001	<0.0001					
	pH	7.51	0.0179	–	–		0.0179		
	Interaction	0.03	0.8633	na	na	na	na	na	na
8	Nutrition	134.3	<0.0001	<0.0001					
	pH	2.7	0.1263	–	–		na		
	Interaction	0.73	0.4101	na	na	na	na	na	na
12	Nutrition	61.69	<0.0001	<0.0001					
	pH	10.04	0.0081	–	–		0.0081		
	Interaction	0.72	0.4137	na	na	na	na	na	na
16	Nutrition	39.43	<0.0001	<0.0001					
	pH	6.19	0.0285	–	–		0.0285	0.8375	0.0816
	Interaction	1.7	0.2162	na	na	na	na	na	na
20	Nutrition	39.91	<0.0001	<0.0001					
	pH	0.56	0.4652	–	–		na		
	Interaction	2.40	0.1472	na	na	na	na	na	na

Significant differences ($P < 0.05$) demonstrated by Student's *t* tests are indicated in bold. *P*-values in merged cells that combine high food (HF) vs. low food (LF) or pH 8.0 vs. pH 7.6 columns correspond to *post hoc* comparisons of means for all HF or LF or pH 8.0 or pH 7.6 data. "na" indicates *post hoc* comparisons that could not be made because of lack of a significant corresponding effect.

Table A5

Experiment 2: linear mixed model assessing juvenile shell length of Crepidula fornicata as a function of time post-metamorphosis, larval pH, and juvenile pH

Fixed effect	df (test, error)	F-ratio	P-value
Days post-metamorphosis	1, 60	368.14	< 0.0001
Larval pH	1, 28	3.65	0.0665
Days post-metamorphosis × larval pH	1, 60	6.07	0.0166
Juvenile pH	1, 28	23.32	< 0.0001
Days post-metamorphosis × juvenile pH	1, 60	126.72	< 0.0001
Larval pH × juvenile pH	1, 28	4.80	0.0369
Days post-metamorphosis × larval pH × juvenile pH	1, 60	13.47	0.0005

The identity of each juvenile culture was included within the model as a random effect. Significant effects ($P < 0.05$) are indicated in bold.

Table A6

Experiment 3: linear mixed model assessing juvenile shell length of Crepidula fornicata as a function of time post-metamorphosis, larval pH, juvenile nutrition, and juvenile pH

Fixed effect	df (test, error)	F-ratio	P-value
Days post-metamorphosis	1, 275	5881.56	< 0.0001
Larval pH	1, 86	23.37	< 0.0001
Days post-metamorphosis × larval pH	1, 275	29.11	< 0.0001
Juvenile food	1, 86	277.74	< 0.0001
Days post-metamorphosis × juvenile food	1, 275	713.80	< 0.0001
Larval pH × juvenile food	1, 86	2.60	0.1113
Days post-metamorphosis × larval pH × juvenile food	1, 275	5.05	0.0255
Juvenile pH	1, 86	3.57	0.0621
Days post-metamorphosis × juvenile pH	1, 275	0.01	0.9531
Larval pH × juvenile pH	1, 86	0.51	0.4758
Days post-metamorphosis × larval pH × juvenile pH	1, 275	0.03	0.8723
Juvenile food × juvenile pH	1, 86	0.08	0.7725
Days post-metamorphosis × juvenile food × juvenile pH	1, 275	0.01	0.9499
Larval pH × juvenile food × juvenile pH	1, 86	0.99	0.3220

The identity of each juvenile culture was included within the model as a random effect. Significant effects ($P < 0.05$) are indicated in bold.

Table A7

Experiment 4: linear mixed model assessing juvenile shell length of Crepidula fornicata as a function of time post-metamorphosis, larval pH, and larval nutrition

Fixed effect	df (test, error)	F-ratio	P-value
Days post-metamorphosis	1, 156	3949.55	< 0.0001
Larval food	1, 36	8.98	0.0049
Days post-metamorphosis × larval food	1, 156	43.86	< 0.0001
Larval pH	1, 36	2.94	0.0950
Days post-metamorphosis × larval pH	1, 156	0.51	0.4765
Larval food × larval pH	1, 36	1.49	0.2304
Larval pH × larval food × days post-metamorphosis	1, 156	5.40	0.0214

The identity of each juvenile culture was included within the model as a random effect. Significant effects ($P < 0.05$) are indicated in bold.